



Makes up to

8 gallons

Concentrate

Kills: Poison ivy, Clover, Ground ivy, Dandelion, Wild Violet and many other listed weeds

Use on Residential

ACTIVE INGREDIENTS: Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*

Dimethylamine Salt of Dicamba (3, 6-Dichloro-o-Anisic Acid)*** . . 1.35%

By Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid

EPA Est. No. 4-NY-1 EPA Reg. No. 228-334-3772

Keep Out Of Reach Of Children
WARNING-AVISO
See Book Panal for First All and
Net Contents 16 FL OZ (473 ML)

EM687889

SUPER BRUSH & WEED KILLER CONCENTRATE

- KILLS CHICKWEED, CLOVER, COCKLEBUR, KNOTWEED, OXALIS, PLANTAIN, THISTLE AND MANY OTHER SPECIES OF BROADLEAF WEEDS.
- ONE PINT COVERS 5,000 SQ. FT.
- CONTAINS MCPA, TRICLOPYR AND DICAMBA

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call 1-800-858-7378

Store and transport in an upright position.

©Bonide Products, Inc. All Rights Reserved

EPA Est. No. 4-NY-1 EPA Reg. No. 228-334-3772

Distributed by

Shenandoah, lowa 51603 Pint





.loh 64801

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Use strictly in accordance with label precautionary statements and directions.

This product is for use on residential turf. IMPORTANT: For use on Bluegrass, Fescue, Rye, Bent, Bermuda, Bahia and Zoysia Lawns. Avoid application to St. Augustine or centipede lawns unless injury can be tolerated. Do not spray Carpet grass, Dichondra or desirable clovers. May cause temporary yellowing of some Bermudagrass turf. Do not exceed specified dosages for any area. Be particularly careful applying within the drip line of trees and other ornamental species. Avoid contact with exposed feeder roots of ornamentals and trees.

Do not apply more than 1.5 lb ae/acre per year. Do not apply more than 2 broadcast applications per year with a minimum retreatment interval of 21 days.

APPLICATION RESTRICTIONS

Entry Restrictions: Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to ente

LAWN WEED CONTROL

When used as directed kills broadleaf weeds in lawns - roots and all. Kills weeds - not lawn grasses.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 pints in 4 to 50 gallons of water per 10,000 square feet (3.2 fluid ounces in 2.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

PG 2

MIXING INSTRUCTIONS (Amount to use)

For Northern Lawns

· Fescues, Kentucky bluegrass, perennial rye & zoysia

When using a tank sprayer: Add 4 fl oz per 1 gallon of water for each 1,000 sq. ft.

For Southern Lawns

Bermuda, bahia, & bent (Do not use on St. Augustine or centipede lawns):

When using a tank sprayer: Add 2 fl oz per gallon of water for each 1,000 sq. ft.

FOR BEST RESULTS

Before You Apply:

Do not mow for 1 to 2 days before or after application.

If soil is dry, water before application.

To treat entire lawn (broadcast application): Use a tank sprayer or hose-end sprayer. Measure lawn. Calculate square feet by multiplying length times width. Spray evenly over measured area.

HOW TO APPLY

To spot treat individual lawn weeds: Use a tank sprayer and spray to wet weeds. To reduce spray drift, adjust sprayer nozzle to deliver a coarse spray.

WHEN TO APPLY

- · Spray when weeds are actively growing.
- · Spray when temperature is below 90° F.
- Spray when air is calm to avoid drift to vegetables, flowers, ornamental plants, trees, shrubs and other desirable plants.
- Do not apply to newly seeded grasses until well-established (they have been mowed 3 times).
- Rain or watering 1 hour after application will not wash away effectiveness.
- Reseed no sooner than 3 weeks after application.

PG 3

BRUSH CONTROL

Kills Brush And Other Tough Weeds - Poison Ivy, Poison Oak, Kudzu, Wild Blackberries and over 240 other tough weeds.

WHERE TO USE

This product may be used around homes, cabins and other buildings and along fences. Amount to Use - 4 fluid ounces per gallon of water per 1.000 square feet.

MIXING INSTRUCTIONS

Amount to Use: 4 fluid ounces per gallon of water per 1,000 square feet (1 tablespoon=3 teaspoons, 1 fluid ounce=2 tablespoons).

When using this product: Set dial to 4 ounces. Pour 4 ounces of this product directly into sprayer jar. DO NOT ADD WATER. Spray evenly over 1,000 square feet. After spraying, unused product should be poured back into its original container.

HOW TO APPLY

- Use coarse spray pattern to reduce drift to desirable plants.
- Spray weeds to thoroughly wet foliage, stems and trunk.
 Do not spray on or around plants or grasses you like, they may die too.
- Do not spray on or around plants or grasses you like, they may die too
- If weeds are growing around desirable plants, shield them from drift with a sheet of cardboard or plastic.
- If desirable plants are accidentally sprayed, rinse off immediately with water.

TO KILL VINES

- If weeds are growing up poles, fences or tree trunks with mature bark, spray to thoroughly cover weeds up to a height of 3 to 4 feet. Tree bark is mature if you scratch the bark with your finger and do not find any green tissue.
- If weeds are climbing shrubs or green barked trees, cut weeds at base and treat as directed for stumps.

TO KILL STUMPS

- Apply product undiluted with a paintbrush to the entire freshly cut stump, including the cut surface, remaining trunk and exposed roots.
- Do not reuse paintbrush. Wrap paintbrush in newspaper and put in trash.

WHEN TO APPLY

- Spray when weeds are actively growing.
- Spray when air is calm to avoid drift to vegetables, flowers, ornamental plants, trees, shrubs and other desirable plants.

Note: Contact with poison ivy or poison oak any time of the year can cause an allergic reaction. Handle dead plants with rubber gloves. Dispose of plants and rubber gloves in tightly sealed garbage bags.

Do not allow people (other than applicator) or pets on treatment area during application. People and pets may enter treated area after spray has dried.

WEEDS

Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Biden, Bindweed, Bird vetch, Bitterweed, Bilter wintercress, Blackeyed Susan, Black medic, Blackmustard, Black-seed plantain, Blessed thistle, Blue leftuce, Blue vervain, Box elder, Bracted plantain, Brasbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebead, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dopánine, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Filixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf

WEEDS Cont.

drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mouseear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed, Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstern filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virgina creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'-clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woodsorrel, Woolly croton, Woolly morningglory, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed and other broadleaf weeds.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: To be stored in original container and placed in an area inaccessible to children.

STORAGE AND DISPOSAL Cont.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

FIRST AID	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomitting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for medical emergency information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING-AVISO: Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, or on clothing. Avoid contact with skin. This product may cause skin sensitization reactions in some people. Wear long-sleeved shirt and long pants, socks and shoes, protective eyewear (goggles or faceshield), and chemical-resistant cloves made of any waterproof material.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

 FNVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

WARRANTY

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ABE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTIES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FINNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMIEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.