

Super Eight™

Insect Control



Termite, Carpenter Ant & Borer Spray

For use outdoors,
around buildings, homes, in
small flower and vegetable
gardens, and on fruit,
nut and ornamental trees

Kills ash borer, dog wood
borer, leafminers and
40 other listed insects



Concentrate
Makes up to
40 gallons!

Keep Out Of Reach Of Children

WARNING (See Back Panel for Additional
Precautionary Statements)

Net Contents 16 FL OZ (473 ML)

72320815

ACTIVE INGREDIENT:

Permethrin . . . (CAS No. 52645-53-1) . . . 13.30%w/w

OTHER INGREDIENTS*86.70%

TOTAL 100.00%

*Contains petroleum distillate

EPA Est. No. 4-NY-1 EPA Reg. No. 4-349-3772

FOR TERMITES: OUTDOOR, POST-CONSTRUCTION,
RESIDENTIAL USE ONLY. DO NOT USE INSIDE DWELLINGS.
For use around existing structures,
not for pre-construction use.

EM682914

EARL MAY **SUPER EIGHT**
**INSECT CONTROL TERMITE,
CARPENTER ANT & BORER SPRAY**

To Control Listed: Ants, Japanese Beetles, Grubs, Mole Crickets, Potato Flea Beetles, Cutworms, Armyworms, Plum Curculio, Pear Psylla, Aphids, Elm Leaf Beetle, Scales, Spider Mites, Mealybugs, Thrips, Whiteflies.

Use on Listed: Lawns, Vegetables, Fruits & Nuts, Flowers & Trees.

Mixture ratios vary. Read and use according to label directions.

EPA Est. No. 4-NY-1

EPA Reg. No. 4-349-3772

Buyers Guarantee Limited to Label Claims

©Bonide Products, Inc. All Rights Reserved

Distributed by
EARL MAY SEED & NURSERY
Shenandoah, Iowa 51603



Pint



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
General Precautions and Restrictions

For use on home lawns, flower and vegetable gardens, fruit, nut and ornamental trees. Do not allow adults, children or pets to enter the treated area until sprays have dried. Do not apply this product in a manner that will contact adults, children or pets, either directly or through drift. Remove pets, birds, and cover fish aquariums before use. Do not apply as a broadcast treatment to indoor surfaces at residential sites including nurseries, day care centers, schools, hospitals and nursing homes.

Not for use on plants being grown for sale or other commercial use or for commercial seed production or for research purposes. Do not water the treated area to the point of run-off. Do not make outdoor applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatments to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- (3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. Not for use in outdoor residential misting systems.

DIRECTIONS FOR USE ON VEGETABLES: For backyard vegetable gardens only. Prepare spray mix as directed in following chart. Use 1 qt. of finished spray per 110 square feet (or 1 gal./440 sq. ft.) of surface area.

HOW TO MIX AND APPLY

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Repeat applications as needed or as noted in the rate chart.

HOW TO MIX AND APPLY Cont.

Step #2: Use clean sprayer. Carefully measure and mix the amount of product and water as suggested in the rate chart. Do not use kitchen utensils for measurements.

Step #3: Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use.

1 fl. oz - 6 tsp. or 2 Tbs.

8 fl. oz - ½ pt. or 1 cup

1 fl. oz. in 3 gallons of water = 2 tsp. in 1 gallon of water or 8 fl. oz. in 24 gallons.

VEGETABLES

Asparagus (Pre-harvest) - To control: Cutworm and Asparagus Beetle. Apply for uniform coverage as needed. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest. Use at a rate of 1 tsp. per 5 qts. of water.

Asparagus (Post-harvest) - To control: Larval stages of Asparagus Beetle, Tarnished Plant Bug, Lygus Bugs, and Adult Japanese Beetle. Treat ferns or bush growth after spear harvest when insects are present. Use at a rate of 1 tsp. per 5 qts. of water.

Celery - To control: Vegetable Leafminer, Cabbage Looper, Black and Granulate Cutworm; Fall, Beet and Southern Armyworm. Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season. Use at a rate of 1 ½ tsp. per 4 qts. of water.

Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower - To control: Beet Armyworm, Diamond Back Moth, Imported Cabbageworm, Cabbage Aphid and Cabbage Looper. Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli, and 10 applications to Cabbage. Use at a rate of 1 tsp per 5 qts. of water.

Eggplant - To control: Colorado Potato Beetle, Vegetable Leafminer. Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest. Use at a rate of 1 ½ tsp. per 4 qts. of water.

Horseradish (Foliar application) - To control: Imported Crucifer Weevil. Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest. Use at a rate of 1 ½ tsp per 4 qts. of water.

VEGETABLES Cont.

Head Lettuce only (Do not use on Leaf Lettuce) - To control: Cabbage Looper, Alfalfa Looper, Beet Armyworm, Corn Earworm and Tobacco Budworm. Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest. Use at a rate of 1½ tsp. per 4 qts. of water.

Potatoes - To control: Beet Armyworm, Colorado Potato Beetle, Potato Leafhopper, Potato Flea Beetle, Aster Leafhopper, Potato Aphid, Potato Tuberworm, Cabbage Looper, Cutworms, European Corn Borer, Potato Psyllid, Tarnished Plant Bug. Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest. Use at a rate of 1½ tsp per 4 qts. of water.

Peppers - To control: Vegetable Leafminer, Cabbage Looper, Flea Beetle, Corn Earworm, Pepper Weevil and European Corn Borer. Apply as needed, but not more than 8 sprays per season. Can be applied up to 3 days prior to harvest. Use at a rate of 1½ tsp. per 4 qts. of water.

Spinach - To control: Beet Armyworm, Corn Earworm, Cabbage Looper, Cutworms, European Corn Borer, Green Cloverworm and Leafhoppers. Apply for uniform coverage when needed, but not more than 7 applications per season. Can be applied up to 7 days prior to harvest. Use at a rate of 1 tsp. per 5 qts. of water.

Sweet Corn - To control: Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm. Apply every 5 days and when needed but not more than 6 applications per season. Can be applied up to 1 day prior to harvest. Use at a rate of 1½ tsp. per 4 qts. of water.

TREES (FRUIT OR NUT)

Almonds - To control: Navel Orange Worm, Peach Tree Borer. Apply when insects appear. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest. Use at a rate of 2¾ tsp. per 5 qts. of water.

Apples - To control: Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique-banded Leafroller and Green Fruitworm. Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall. Use at a rate of 2¾ tsp. per 5 qts. of water.

TREES (FRUIT OR NUT) Cont.

Peaches - To control: Lesser Peachtree Borer, Peach Twig Borer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth. Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season. Use at a rate of 2¼ tsp. per 5 qts. of water.

Pears (dormant through delayed dormant) - To control: Pear Psylla. Apply during the dormant through delayed dormant growth period only. Add if desired, 8 oz. of Bonide All Season's Oil to each gallon of Earl May's Super Eight Insect Control Concentrate and use spray. Do not make more than 2 applications per season. Use at a rate of 2¼ tsp. per 5 qts. of water.

Pears (summer) - To control: Pear Psylla, Green Fruitworm and Codling Moth. Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest. Use at a rate of 2¼ tsp. per 5 qts. of water.

FOR ORNAMENTAL TREES GROWN AROUND THE HOME

Use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: Boxelder bug, Clearwing moth borer, Ash borer, Banded ash clearwing borer, Dogwood borer, Lesser peachtree borer, Lilac borer, Oak borer, Peachtree borer, Rhododendron borer, Coleopteran borer, Bronze birch borer, Flatheaded appletree borer, Pine Moths, Pine Needle Miners, Pine Beetles and Needle Scales on pine, Budworms on conifers, Oakworms on oak, Fall Webworms on cherry, elm and honeysuckle, Aphids on tulip poplar, Citrus Black Flies on non-bearing citrus, Tussock moths on Douglas and grand firs, Leafminers on birch, Elm Leaf Beetle and Elm Spanworms on elms. Mix 2¼ tsp. per 5 qts. of water.

TO KILL TENT CATERPILLARS

Application should be made when caterpillars are young and tents are first noticed. For results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Mix 2 fl. oz. of this product in one gallon of water. Wet tents with spray on all sides. A few caterpillars may be seen escaping from tent; however, they should die within several hours.

FOR USE ON HOME LAWNS

For control of ants (including fire ants), armyworms, chinch bugs, crickets, cutworms, Japanese beetle grubs, grasshoppers, mole crickets and sod webworms, consult the following chart:

Pests	Dilution
Armyworms, Cutworms, Chinch Bugs, Japanese Beetle Grubs, Mole Crickets, Sod Webworm	5 tsp./10 gal. water/1000 sq. ft.
Crickets, Grasshoppers, Ants (Including Fire Ants)	2 tsp./3 gal. water/1000 sq. ft.

Lawn should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed above. For heavy infestations, repeat application after two weeks.

Also for control of cutworms on dichondra and Irish moss. Use as directed above for lawns.

Keep children and pets off treated areas until lawn has dried following the application.

FOR USE ON ORNAMENTAL PLANTS OUTDOORS

Contact and Residual Spray - Use sufficient spray to obtain full coverage of all foliage. Adjust spray volume used to plant size.

To use, mix 1 teaspoon of Earl May's Super Eight Insect Control Concentrate with one gallon of water.

Spray in the late afternoon or evening, when the temperature ranges from 50° to 75° F and when there is little or no wind. Spray at the first indication of insect pests. On those plants that will flower during the season, the spray schedule should be completed before blossom formation. Repeat as necessary; use intervals of 4 to 8 days. Application can be made up to the day of collection.

Spray as directed to control aphids, cicadas (locust), corn earworms, fall cankerworms, inchworms, leafminers, leafrollers, locust, mealybugs, bagworms, spider mites, exposed thrips and whiteflies on the following plants: Ageratum, Arizona Cypress, Aster, Azalea, Begonia, Coleus, Common Ninebark, Common Snowberry, Elm,

FOR USE ON ORNAMENTAL PLANTS OUTDOORS Cont.

Exacum, Euonymous, English Ivy, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Taxus, Wax Plants, Weeping Fig, Zinnia.

OUTDOOR PEST CONTROL

For contact and residual spray, mix with water in the following manner:

Finished Spray	Insect Control (Fluid Ounces)	Water
0.25%	2 2/3 (5 1/3 Tbs.)	1 gallon
0.25%	2/3 (4 tsps.)	1 quart
0.50%	5 1/2 (10 2/3 Tbs.)	1 gallon
0.50%	1 1/3 (8 tsps.)	1 quart

For heavy infestations, use the 0.50% dilution. For lighter infestations and for maintenance purposes, use the 0.25% dilution. One gallon of finished spray is sufficient to treat 1580 linear feet with a 6 inch spray band.

PESTS ON OUTSIDE SURFACES OF BUILDING

Use this product to control ants, Asian lady beetle, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 3 feet where pests are and may find entrance. Also, apply as a spot treatment to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FLYING MOTHS, OUTDOORS: For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

ANTS, DEER AND LONE STAR TICKS, CRICKETS, FLEAS, CICADAS (LOCUST), AND EARWIGS, OUTDOORS: For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

WASPS, HORNETS, YELLOW JACKETS, BEES (including CARPENTER BEES), OUTDOORS: Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

TO CONTROL FIRE ANTS: Dilute 2 tablespoons of this product in 1 gallon of water. Apply 1 gallon of solution as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon per mound is still required. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

SUBTERRANEAN TERMITES: DO NOT USE INSIDE DWELLINGS. For use around existing structure, not for pre-construction use. For post-construction treatment around residential buildings. **Mix 5 fl. oz. of Earl May's Super Eight Insect Control Concentrate per 1 gallon of water (makes a 0.5% water emulsion).** Apply as specified below. One treatment should last up to 5 years if treated soil is left undisturbed.

Kills and controls subterranean termites. Chemicals for soil treatment are used to establish a barrier against termites. The chemical emulsion must be adequately dispersed in the soil to provide a barrier between the wood in the structure and the termite colonies in the soil. Barriers may be established in the soil on the outside perimeter of the house by trenching using a rate of 4 gallons of emulsion per 10 linear feet per foot of depth. A complete termite barrier may require treatment with another EPA registered product to the inside perimeter of the foundation and to other interior critical areas. For these situations, contact a licensed pest control operator for treatment. Do not use under slabs, in crawl spaces, or indoors. If termites are found within the house or structure, contact a licensed pest control operator for treatment.

Remove all wood debris and wood forms. Around piers, pipes, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. For brick and block foundations, dig the trench about 12 inches deep. If footing is more than 12 inches deep, make holes with a crowbar, pipe or a rod about 1 foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. NEVER DIG BELOW THE TOP OF THE FOOTING. Apply the prepared emulsion (or solution) in the trench and to the backfill soil at a rate of 4 gallons per 10 linear feet per foot of depth. Cover treated soil with a thin layer of untreated soil. For raised porches, terraces and entrance slabs outdoors, drill holes at one foot intervals, 6 inches from the foundation, and pour one-half gallon of emulsion (or solution) into each hole. Refill holes.

Be sure to treat the interiors of accessible hollow block foundations at same rate as above. Do not use in structural voids insulated with rigid foam. Be careful to have adequate ventilation when treating areas under structure. For severe infestations or treatment under slabs with heat ducts or radiating heat pipes, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service.

NOTE: Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water.

Use anti-backflow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware or plumbing supply stores. Use of anti-backflow equipment will help ensure that the user of this chemical will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Do not treat structures that contain cisterns, wells, French drains, or sump systems. Soil in the vicinity of wells, cisterns or ponds should not be treated. For those situations, contact the local Cooperative Extension Service or consult a commercial pest control service. Do not treat soil that is water saturated or frozen or in any conditions where runoff or movement from the treatment area (site) is likely to occur.

WHEN TO RE-APPLY

Reapplication for subterranean termites should only be made when there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas. Annual reapplication of the entire premises must be avoided.

CARPENTER ANTS, CARPENTER BEES & WOOD-INFESTING BEETLES: For localized treatment in and around residential buildings. **Mix 5 fl. oz. of Earl May's Super Eight Insect Control Concentrate in 1 gallon of water and apply as directed.** Allow spray to dry before allowing people or pets to occupy treated areas.

Carpenter Ants: Adjust sprayer to deliver a coarse spray. Apply mixture around doors, windows and other points of entry. Spray into cracks and crevices where ants or their nests are suspected.

Carpenter Bees & Wood-Infesting Beetles (Old House Borer and Powder Post): To protect wood and control damage by these wood-infesting insects, thoroughly apply mixture to wood using a coarse spray or using a paintbrush. Do not allow runoff to occur.

STORAGE & DISPOSAL

Pesticide Storage: Do not store at temperatures below 40°F (4.5°C). Store product in original container in a locked storage area, inaccessible to children and pets. Store and transport in upright position.

Pesticide Disposal & Container Handling: *If empty* - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. *If partly filled* - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Note to Physician: Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or safety glasses. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Flashpoint minimum of 105°F (49°C).

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

To the extent consistent with applicable law, buyer assumes all risks for use of this product when not used according to label directions or established safe practice.